

## Literature Critique Criteria for Systematic reviews and meta-analyses - tabular form

Criterion	Green	Yellow	Red	Comments
The study is in fact identified as a systematic review or meta-analysis	“Systematic review,” “meta-analysis,” or both, are in the title of the article, and the abstract supports the design in the title	The title is ambiguous, but the abstract shows that the authors did a systematic review	The article is a narrative review or an overview, or is done by a single author	“Systematic review” and “meta-analysis” are generally recognized terms for a specific type of original research; narrative reviews are subject to biases which systematic reviews and meta-analyses methodically control for
Objectives of the systematic review or meta-analysis	Clearly stated in terms of PICOS: <b>P</b> atient population (disease, age, setting), <b>I</b> ntervention (dose, frequency, etc), <b>C</b> omparator (control group interventions), <b>O</b> utcome (morbidity, mortality, symptoms, function), and <b>S</b> tudy design (randomized trials only, broader design criteria)	PICOS elements all reported, but some ambiguity in some elements (e.g., Comparator described as “standard care” or “usual care” without further description)	One or more PICOS element missing or uninterpretable	The question being addressed should be clear from the abstract; it may be narrow or broad, but the scope and potential applicability should be well defined
Characteristics of eligible	In addition to PICOS, study	Ambiguity exists for some	Eligibility of studies is	

Criterion	Green	Yellow	Red	Comments
studies	characteristics defined in terms of restrictions for inclusion (e.g., minimum length of follow-up, whether co-interventions are included), and scope of reports (language, years of publication, unpublished material)	of the characteristics of eligible studies	unclear, and scope of reports is not specified	
Information sources	Multiple information sources are clearly specified: databases (PubMed, Ovid, EMBASE, Cochrane, Web of Science), hand searches of tables of contents of relevant journals, meeting abstracts, reference lists, contacts with authors, manufacturers, trial registries)	Search limited to published material from two or more sources, without additional searching of registries or contact with authors	Search limited to a single information source (e.g., PubMed only)	It is desirable to search multiple databases beyond PubMed, but there is little evidence that data sources beyond PubMed lead to different conclusions in meta-analyses; this criterion, if not met, is not a fatal flaw in a systematic review or meta-analysis
Search strategy	Full electronic search strategy for at least one major database, with dates (e.g.,	Databases and search terms are given, but there is some ambiguity in	Databases and search terms are too broad and vague to permit	Often given in an appendix to the article or in an online supplement, the

Criterion	Green	Yellow	Red	Comments
	PubMed 1970-October 2009), limits, combinations of search terms, such that it can be replicated by the reader	the strategy (e.g., PubMed “through 2007”), and replication by the reader would be difficult	replication by the reader	strategy should be readily accessible
Study selection	Specification of which criteria determine eligibility for inclusion (e.g., randomization to specified interventions, which outcomes were required to be reported) and for quality that is consistent with our criteria (e.g., allocation concealment, intention-to-treat analysis, blinding) with at least two reviewers identified by initials; inter-rater agreement and methods of resolving disagreement are specified; a flow diagram enumerates articles retrieved from search, articles excluded after screening, and articles	Two or more reviewers screen articles for inclusion, but there is some ambiguity in the criteria for inclusion or for inter-rater agreement and methods of resolving disagreement; flow diagram is lacking articles that do not qualify under our criteria as adequate evidence will not be used to establish evidence statements.	Only one reviewer selects studies; criteria are vague or do not clearly match our criteria for RCT.	Quality assessment should focus on risk of bias; scoring of articles for quality is not necessary and may be misleading. There is no standard process for selecting studies, but the process used by the reviewers should be clear enough to allow the reader to determine which studies might meet the test of inclusion

Criterion	Green	Yellow	Red	Comments
	included for meta-analysis			
Outcomes for analysis	Meta-analysis is restricted to pre-specified primary and secondary outcomes, and exploratory (hypothesis-generating) analyses in the source literature are excluded from meta-analysis	Meta-analysis combines pre-specified primary and secondary outcomes in the source literature with exploratory analyses in the same literature, but assigns exploratory analyses a lower weight	Meta-analysis treats exploratory analyses in source literature on an equal basis with the pre-specified primary and secondary analyses	Exploratory analyses are too likely to be reported when they arise from the play of chance, and should not be included in any meta-analysis of the same outcomes; their inclusion is likely to bias the meta-analysis
Summary measures for meta-analysis with or without pooled Number Needed to Treat (NNT)	Principal summary measures (relative risk, risk difference, odds ratio, difference in means, hazard ratio) are specified and appropriate to the outcome measure; if numbers needed to treat (NNT) are reported, there is a fixed event rate in the control groups for the studies being combined	Risk ratios or odds ratios are reported, and NNT is not reported if there is a difference in the control group event rates across the different studies	Risk ratios or odds ratios are reported, but NNT is reported even when there is a difference in control group event rates across the different studies (the underlying baseline risks are not equal)	Relative risks and odds ratios are generally more stable for summary measures than risk differences; pooled NNT is misleading if the control group event rate (the baseline risk) is different across studies, even if the risk ratio is the same
Meta-analysis presentation	Results of meta-analysis are presented as an estimated summary effect	Estimated summary effect with confidence interval, with an estimate of	Summary effect measure with confidence interval, but heterogeneity	No hard and fast rule dictates the choice of model, but

Criterion	Green	Yellow	Red	Comments
	<p>(with confidence interval) across all included studies, displaying a forest plot with weights and confidence intervals for the included studies; a measure of heterogeneity is presented (e.g., <math>I^2</math>); the choice of fixed effect or random effects model is explained, and, if there is significant heterogeneity, there is an attempt to examine possible sources of heterogeneity</p>	<p>heterogeneity, and an explanation of the choice of fixed or random effects model; however, an examination of sources of heterogeneity is lacking</p>	<p>measures and examinations are lacking</p>	<p>because a fixed effect model assumes a single common effect size across studies, there should be a discussion of why it is appropriate for the included studies</p>

Cumulative Trauma Conditions - 2016